

Factorizations in universal enveloping algebras of three dimensional Lie algebras and generalizations

STEPHEN BERMAN¹, JUN MORITA² and YOJI YOSHII³

- 1) Department of Mathematics, University of Saskatchewan
Saskatoon, Saskatchewan, Canada S7N 5E6
berman@snoopy.usask.ca
- 2) Institute of Mathematics, University of Tsukuba
Tsukuba, Ibaraki, 305-8571, Japan
morita@math.tsukuba.ac.jp
- 3) Department of Mathematics, University of Alberta
Edmonton, Alberta, Canada, T6G 2G1
yoshii@math.ualberta.ca

This paper is dedicated to Robert V. Moody on the occasion of his 60th birthday

Abstract

We introduce the notion of Lie algebras with plus-minus pairs as well as regular plus-minus pairs. These notions deal with certain factorizations in universal enveloping algebras. We show that many important Lie algebras have such pairs and we classify, and give a full treatment of, the three dimensional Lie algebras with plus-minus pairs.

AMS Classification Numbers 17B05, 17B35, 17B67, 17B70

1 Introduction

All of our algebras will be over a field F of characteristic zero. We begin by recalling the well known fact that if L is a Kac-Moody Lie algebra with the usual Chevalley generators $\{e_i, f_i | 1 \leq i \leq l\}$ satisfying $L = [L, L]$ and l is finite, then every L -module on which the elements $e_i, f_i, 1 \leq i \leq l$ act locally nilpotently is integrable in the sense that the elements $h_i = [e_i, f_i], 1 \leq i \leq l$ are simultaneously diagonalizable. (cf. [10] Ex. 6.31, p. 585, or [11]). In other words, every weakly integrable module for such an algebra is integrable. The usual proof of this fact uses that the three dimensional Lie algebra with basis e_i, f_i, h_i is isomorphic to the Lie algebra sl_2 (so $\mathfrak{g}$ is generated by sl_2 -triples) together with the result which says that if V is any module for sl_2 on which the standard generators e, f of sl_2 act locally nilpotently then the element $h = [e, f]$ is diagonalizable on V . One can see this last fact as follows. We denote by $M(W)$ the maximal integrable submodule

of an sl_2 -module W . In general, $M(W/M(W)) = 0$ for all sl_2 -modules W . If a vector v of an sl_2 -module W satisfies that $fv = 0$, $e^n v \neq 0$ and $e^{n+1}v = 0$, then we obtain $h(e^n v) = n(e^n v)$, since $fe^n = e^n f - ne^{n-1}(h + n - 1)$ and

$$\begin{aligned} h(e^n v) &= (ef - fe)e^n v \\ &= efe^n v \\ &= e(e^n f - ne^{n-1}(h + n - 1))v \\ &= -ne^n(h + n - 1)v \\ &= -n(h - n - 1)e^n v \\ &= -nh(e^n v) + n(n + 1)e^n v. \end{aligned}$$

This implies that $M(V)$ is nontrivial for every nonzero sl_2 -module V on which the elements e and f are locally nilpotent operators. Therefore, $M(V) = V$ for such an sl_2 -module V , that is, V is integrable.

We want to indicate another approach to the above fact about sl_2 -modules which uses a factorization in the universal enveloping algebra. This method appears to be new and was the starting point of this paper. For any Lie algebra $\mathfrak{g}$ we let $U(\mathfrak{g})$ denote its universal enveloping algebra. Then one knows that for the algebra $sl_2 = Fe \oplus Fh \oplus Ff$ we have the factorization

$$U(sl_2) = U(Fe)U(Ff)U(Fh). \quad (1)$$

Using this it is easy to see that if V is an sl_2 -module on which both e and f act locally nilpotently then any vector $v \in V$ generates a finite dimensional submodule. Thus h acts semisimply on this submodule and so we obtain h acts semisimply on V . Also, the proof of the factorization (1) is quite straightforward and follows easily from the following formula in $U(sl_2)$.

$$f(e^i f^j e^k) = \frac{j-i+1}{j+1} e^i f^{j+1} e^k + \frac{i}{j+1} e^{i-1} f^{j+1} e^{k+1} + i(j-i+1) e^{i-1} f^j e^k, \quad (2)$$

for all $i > 0, j, k \geq 0$. This formula can be established using the following: for any $k \geq 0$ we have

$$\begin{aligned} (A_k) \quad f e f^k &= \frac{k}{k+1} e f^{k+1} + \frac{1}{k+1} f^{k+1} e + k f^k \\ (B_k) \quad f^k e f &= \frac{1}{k+1} e f^{k+1} + \frac{k}{k+1} f^{k+1} e + k f^k. \end{aligned}$$

$f e^i = e^i f - i e^{i-1}(h + i - 1)$ for $i \geq 1$ and induction.

Next let $\mathfrak{H}$ be the three dimensional Heisenberg Lie algebra with a basis x, y, z satisfying $[x, y] = z$, $[x, z] = [y, z] = 0$. Then, in $U(\mathfrak{H})$, we obtain the following factorization

$$U(\mathfrak{H}) = U(Fx)U(Fy)U(Fz). \quad (3)$$

The proof of this is much like the sl_2 case. It follows easily from the following formula in $U(\mathfrak{H})$

$$y(x^i y^j x^k) = \frac{j-i+1}{j+1} x^i y^{j+1} x^k + \frac{i}{j+1} x^{i-1} y^{j+1} x^{k+1}, \quad (4)$$

for all $i > 0, j, k \geq 0$. Note that (4) is proved by establishing for any $k \geq 0$ we have

$$\begin{aligned} (A'_k) \quad y x y^k &= \frac{k}{k+1} x y^{k+1} + \frac{1}{k+1} y^{k+1} x \\ (B'_k) \quad y^k x y &= \frac{1}{k+1} x y^{k+1} + \frac{k}{k+1} y^{k+1} x, \end{aligned}$$

which in turn is proved using $yx^i = x^iy - ix^{i-1}z$ for $i \geq 1$ and induction. Thus, the picture is similar for both algebras sl_2 and $\mathfrak{H}$ in that they both have a pair of subalgebras P, M satisfying $P + M$ is not the whole algebra and $U(P)U(M)U(P)$ is the whole enveloping algebra. This prompts the following definition which singles out those Lie algebras having this type of factorization in their universal enveloping algebras.

Definition 1.1 (i.) A Lie algebra L is said to have a **plus-minus pair** if it has two subalgebras P, M satisfying $P + M \neq L$ and

$$U(L) = U(P)U(M)U(P).$$

In this case we say L has a plus-minus pair (P, M) .

(ii.) Let (P, M) be a plus-minus pair of L . We say this is a **regular plus-minus pair** if $P \cap M = (0)$ and there is an automorphism σ of L of order two satisfying $\sigma(P) = M$. Note that in this case we then have $U(L) = U(P)U(M)U(P) = U(M)U(P)U(M)$.

It is clear that both Lie algebras sl_2 and $\mathfrak{H}$ have regular plus-minus pairs. Moreover if L is any three dimensional Lie algebra with a plus-minus pair (P, M) then each of P and M must be one dimensional. Indeed, $P + M$ cannot be 3 dimensional as $P + M \neq L$. Thus, $P + M$ is two dimensional and so if one of P, M is 2 dimensional then $P + M$ is a subalgebra of L and so $U(P + M) \neq U(L)$ but $U(P)U(M)U(P) \subseteq U(P + M)$ so (P, M) cannot be a plus-minus pair. Letting $P = Fx, M = Fy$ we have that

$$U(L) = \sum_{i,j,k \geq 0} Fx^i y^j x^k. \quad (5)$$

Moreover, the following result, which extends the situation discussed in the sl_2 case, is quite clear. Let L be a three dimensional Lie algebra with a plus-minus pair (P, M) where

$P = Fx, M = Fy$. Let V be any L -module on which the action of the elements x, y is locally finite. Then any finitely generated submodule of V is finite dimensional. Thus, one is led to ask just which three dimensional Lie algebras have plus-minus pairs.

In Section 2 we will extend the methods used in the proofs for the sl_2 and $\mathfrak{H}$ cases above and show that any three dimensional Lie algebra which is generated by two elements has a plus-minus pair. Then, using results on three dimensional Lie algebras from [6] we will see that there are only two isomorphism classes of the three dimensional Lie algebras which do not have plus-minus pairs. We also go on to study, when the base field F is algebraically closed, which of these algebras have regular plus-minus pairs and are able to give a complete list of these. In the third and final section of this paper we go on to investigate plus-minus pairs, or similar factorizations, in the universal enveloping algebras, of Borchers Lie algebras as well as in some $\mathbf{Z}^n$ -graded Lie algebras which satisfy some extra conditions.

2 Three dimensional case

In this section we begin by showing a three dimensional Lie algebra has a plus-minus pair if and only if it is generated by two elements. We then go on to investigate some special cases as well as regular plus-minus pairs when the base field is algebraically closed.

Throughout we let L be a three dimensional Lie algebra unless mentioned otherwise. If (P, M) is a plus-minus pair for L then we know that each of P, M is one dimensional so we let $P = Fx, M = Fy$. If x and y do not generate L then it must be that $P + M$ is a proper subalgebra of L and so since $U(P)U(M)U(P) \subseteq U(P + M)$ we get a contradiction. Thus L is generated by x and y so is two-generated. We want to establish the converse of the above result. For the moment we let L be an arbitrary Lie algebra and x, y any two elements of L .

We define subspaces U_k for $k \geq 0$ of $U(L)$ by saying

$$U_k = \sum_{0 \leq m \leq k} (Fxy^m + Fy^m x). \quad (6)$$

Notice that $U_0 = Fx$ and that $U_k \subseteq U_{k+1}$ for all $k \geq 0$.

Lemma 2.1 Let L be an arbitrary Lie algebra and x, y any two elements of L . For $k \geq 0$ the following statements hold,

$$\begin{aligned} (A_k) \quad & yxy^k \equiv \frac{k}{k+1}xy^{k+1} + \frac{1}{k+1}y^{k+1}x \pmod{U_k}, \\ (B_k) \quad & y^kxy \equiv \frac{1}{k+1}xy^{k+1} + \frac{k}{k+1}y^{k+1}x \pmod{U_k}, \\ (C_k) \quad & y U_k \subset U_{k+1}, \quad U_k y \subset U_{k+1}. \end{aligned}$$

Proof. We prove this by induction on k noting that for $k = 0$ both (A_0) and (B_0) are clear. Next, we show $(A_0), \dots, (A_k), (B_0), \dots, (B_k)$ imply (C_k) . Indeed, by definition we have that

$$y U_k = \sum_{0 \leq m \leq k} (Fyxy^m + Fy^{m+1}x)$$

and so by $(A_0), \dots, (A_k)$ we get that this is contained in U_{k+1} . Similarly we have

$$U_k y = \sum_{0 \leq m \leq k} (Fxy^{m+1} + Fy^m xy)$$

and so by $(B_0), \dots, (B_k)$ we get this is contained in U_{k+1} . Hence (C_k) holds.

Next we show that $(A_k), (B_k), (C_k)$ imply $(A_{k+1}), (B_{k+1})$. Now $yxy^{k+1} = (yxy^k)y$ so that (A_k) implies that the difference

$$yxy^{k+1} - \left(\frac{k}{k+1}xy^{k+1} + \frac{1}{k+1}y^{k+1}x \right)y \in U_k y.$$

But (C_k) implies that $U_k y \subseteq U_{k+1}$ so we get that

$$yxy^{k+1} \equiv \left(\frac{k}{k+1}xy^{k+1} + \frac{1}{k+1}y^{k+1}x \right)y \pmod{U_{k+1}}.$$

Similarly, using (B_k) and (C_k) we get that

$$y^{k+1}xy \equiv y \left(\frac{1}{k+1}xy^{k+1} + \frac{k}{k+1}y^{k+1}x \right) \pmod{U_{k+1}}.$$

Thus, we finally get

$$yxy^{k+1} \equiv \frac{k}{k+1}xy^{k+2} + \frac{1}{(k+1)^2}yxy^{k+1} + \frac{k}{(k+1)^2}y^{k+2}x \pmod{U_{k+1}}$$

which implies that

$$\frac{k(k+2)}{(k+1)^2}yxy^{k+1} \equiv \frac{k}{k+1}xy^{k+2} + \frac{k}{(k+1)^2}y^{k+2}x \pmod{U_{k+1}}.$$

Therefore, we obtain

$$yxy^{k+1} \equiv \frac{k+1}{k+2}xy^{k+2} + \frac{1}{k+2}y^{k+2}x \pmod{U_{k+1}},$$

and we see that (A_{k+1}) holds.

Using a similiar type of argument we have that

$$\begin{aligned} y^{k+1}xy &= y(y^kxy) \\ &\equiv y\left(\frac{1}{k+1}xy^{k+1} + \frac{k}{k+1}y^{k+1}x\right) \pmod{U_{k+1}} \\ &\equiv \frac{1}{k+1}yxy^{k+1} + \frac{k}{k+1}y^{k+2}x \pmod{U_{k+1}} \\ &\equiv \frac{1}{k+1}\left(\frac{k}{k+1}xy^{k+1} + \frac{1}{k+1}y^{k+1}x\right)y + \frac{k}{k+1}y^{k+2}x \pmod{U_{k+1}} \\ &\equiv \frac{k}{(k+1)^2}xy^{k+2} + \frac{1}{(k+1)^2}y^{k+1}xy + \frac{k}{k+1}y^{k+2}x \pmod{U_{k+1}} \end{aligned}$$

and

$$\frac{k(k+2)}{(k+1)^2}y^{k+1}xy \equiv \frac{k}{(k+1)^2}xy^{k+2} + \frac{k}{k+1}y^{k+2}x \pmod{U_{k+1}}.$$

Therefore, we obtain

$$y^{k+1}xy \equiv \frac{1}{k+2}xy^{k+2} + \frac{k+1}{k+2}y^{k+2}x \pmod{U_{k+1}},$$

and we see that (B_{k+1}) holds. This completes our induction. Q.E.D.

We apply this Lemma to the three dimensional case in our next result.

Theorem 2.2 Let L be a three dimensional Lie algebra. Then L has a plus-minus pair if and only if L is two generated. Moreover, if x and y generate L then (P, M) is a plus-minus pair for L where $P = Fx, M = Fy$.

Proof. We need only show L has a plus-minus pair if L is generated by two elements x, y . Let $z = [x, y]$. Now we want to show $U(L) = \sum_{i,j,k \geq 0} Fx^i y^j x^k$. Put $\mathfrak{X} = \sum_{i,j,k \geq 0} Fx^i y^j x^k \subset U(L)$ and let U_k be defined as above. Clearly $x\mathfrak{X} \subset \mathfrak{X}, \mathfrak{X}x \subset \mathfrak{X}$ and $U_k \subseteq \mathfrak{X}$ for all $k \geq 0$. We claim

$$\begin{aligned} y(x^\ell y^m x^n) &\in \mathfrak{X}, \\ z(x^\ell y^m x^n) &\in \mathfrak{X}. \end{aligned}$$

and show this by induction on ℓ . If $\ell = 0$, then we see $y(y^m x^n) \in \mathfrak{X}$ and using (A_m) we get

$$\begin{aligned} z(y^m x^n) &= (xy - yx)(y^m x^n) \\ &= xy^{m+1}x^n - yxy^m x^n \\ &\in Fxy^{m+1}x^n + (Fxy^{m+1} + Fy^{m+1}x + U_m)x^n \subset \mathfrak{X}. \end{aligned}$$

Let $\ell > 0$. Then, we obtain, using our inductive assumption, that

$$\begin{aligned} y(x^\ell y^m x^n) &= (xy - z)(x^{\ell-1} y^m x^n) \\ &\in x\mathfrak{X} + \mathfrak{X} \subset \mathfrak{X} \end{aligned}$$

and, letting $[z, x] = ax + by + cz$ for $a, b, c \in F$, we also get using our inductive assumption that

$$\begin{aligned} z(x^\ell y^m x^n) &= (xz + ax + by + cz)(x^{\ell-1} y^m x^n) \\ &\in x\mathfrak{X} + \mathfrak{X} + \mathfrak{X} + \mathfrak{X} \subset \mathfrak{X}. \end{aligned}$$

Hence, $y\mathfrak{X} \subset \mathfrak{X}$. Since $\mathfrak{X}$ is a left ideal of $U(L)$ containing 1, we obtain $\mathfrak{X} = U(L)$. Therefore, (P, M) with $P = Fx$ and $M = Fy$ is a plus-minus pair for L . Q.E.D.

If our three dimensional Lie algebra L is abelian it clearly does not have a plus-minus pair. Also, we let $\mathfrak{g}$ be the three dimensional Lie algebra with basis x, y, z satisfying

$$[x, y] = 0, [x, z] = x, [y, z] = y.$$

Then for any elements $a, b, c, \alpha, \beta, \gamma \in F$ we have the very special identity

$$[ax + by + cz, \alpha x + \beta y + \gamma z] = \gamma(ax + by + cz) - c(\alpha x + \beta y + \gamma z).$$

This clearly implies that $\mathfrak{g}$ is not two generated so does not have a plus-minus pair. Our next result shows that these are the only two kinds of three dimensional Lie algebras which do not have plus-minus pairs. In the proof we will make use of the classification of three dimensional algebras presented in Jacobson's book, [6], pg.11-14. We break our argument into 5 subcases (labeled (a.) through (e.)) just as he does. Here L' denotes the derived algebra of L .

Theorem 2.3 Let L be a three dimensional Lie algebra. Then L has a plus-minus pair if and only if L is not abelian and is not isomorphic to the algebra $\mathfrak{g}$ above.

Proof. (a.) If $L' = (0)$ then L is abelian and it clearly has no plus-minus pair.

(b.) Assume L' is one dimensional and in the center, $Z(L)$, of L . Then there is only one Lie algebra satisfying these conditions and it has basis x, y, z satisfying $[x, y] = z, [x, z] = [y, z] = 0$. This is clearly generated by x, y so has a plus-minus pair.

(c.) Assume that L' is one dimensional but not in $Z(L)$. Then L has basis x, y, z satisfying $[x, y] = x, [x, z] = [y, z] = 0$ so L is generated by the two elements $x + z$ and y , hence has a plus-minus pair.

(d.) Suppose that L' is two dimensional. Then, from [6], one has that L' is abelian with basis, say x, y , and there is an element $z \in L$ so that $[x, y] = 0, [x, z] = \alpha x + \beta y, [y, z] = \gamma x + \delta y$ with $\alpha\delta - \beta\gamma \neq 0$. If $\alpha = 0$, then x and $y + z$ generate L since $[x, y + z] = \beta y$ and $\beta \neq 0$ as $\alpha = 0$.

Suppose $\alpha \neq 0$. By replacing z by z/α , we can assume $\alpha = 1$. Hence, we have that $[x, y] = 0, [x, z] = x + \beta y, [y, z] = \gamma x + \delta y$ with $\delta \neq \beta\gamma$. If $\gamma \neq 0$, then $-\gamma x + y$ and z generate L

since $[-\gamma x + y, z] = (\delta - \beta\gamma)y$. Next suppose $\gamma = 0$. Then we have $\delta \neq 0$, and our relations are $[x, y] = 0, [x, z] = x + \beta y, [y, z] = \delta y$. If $\beta \neq 0$, then $x - (\beta/\delta)y$ and z generate L since $[x - (\beta/\delta)y, z] = x$. In the remaining case, our relations are $[x, y] = 0, [x, z] = x, [y, z] = \delta y$ with $\delta \neq 0$. If $\delta \neq 1$, then $x + y$ and z generate L since $[x + y, z] = x + \delta y$. Therefore the only case left is when $\delta = 1$ and then our algebra L is nothing but the algebra $\mathfrak{g}$ above.

(e.) This is the case where $L' = L$. Then Jacobson shows there is a basis x, y, z of L satisfying $[x, y] = z, [x, z] = \alpha y, [y, z] = \beta x$ for some $\alpha, \beta \in F$ with $\alpha\beta \neq 0$. Here we see that x and y generate L . Q.E.D.

The special case when $L = L_{-1} \oplus L_0 \oplus L_1$ is a three graded Lie algebra of dimension three with a plus-minus pair can be discussed without reference to Jacobson's classification. We shall use this in the final section of this work so we briefly go through this here. Put $L_1 = Fx, L_{-1} = Fy, L_0 = Fz$. We can assume first that $[x, y]$ is either 0 or z . Suppose $[x, y] = 0$. If $[x, z] = [y, z] = 0$, then L is abelian so has no plus-minus pair. If $[x, z] = 0$ and $[y, z] \neq 0$, then we can also suppose $[z, y] = y$ and hence, $P = F(x + y)$ and $M = Fz$ give a plus minus pair. If $[x, z] \neq 0$ and $[y, z] = 0$, then we can suppose $[z, x] = x$ and hence, again $P = F(x + y)$ and $M = Fz$ becomes a plus-minus pair. If $[x, z] = ax$ and $[y, z] = by$ with $ab \neq 0$, then we can suppose $a = 1$. In this case, $P = F(x + y)$ and $M = Fz$ give a plus-minus pair when $b \neq 1$. Otherwise we have $[x, z] = x$ and $[y, z] = y$ and there is no

plus-minus pair. Next we suppose $[x, y] = z$. If $[x, z] = [y, z] = 0$, then L is a Heisenberg Lie algebra, and hence, L has a plus-minus pair. If $[x, z] = ax$ and $[y, z] = by$ with $a \neq 0$ or $b \neq 0$, then $0 = [z, z] = [[x, y], z] = [[x, z], y] + [x, [y, z]] = a[x, y] + b[x, y] = (a + b)z$ and $a + b = 0$. Put $x' = x, y' = -2y/a, z' = -2z/a$. Then, $[x', y'] = -2[x, y]/a = -2z/a = z', [z', x'] = -[x', z'] = 2[x, z]/a = 2x = 2x'$ and $[z', y'] = -[y', z'] = -4[y, z]/(a^2) = 4y/a = -2y'$. This means that L is isomorphic to sl_2 . Therefore we obtain the following result which gives a characterization of sl_2 and $\mathfrak{H}$.

Proposition 2.4. Let $L = L_1 \oplus L_0 \oplus L_{-1}$ be a three graded Lie algebra of dimension three with $\dim L_{\pm 1} = \dim L_0 = 1$.

- (1) If L has a plus-minus pair, then L is isomorphic to one of $sl_2, \mathfrak{H}$ and $K(a, b)$, where $K(a, b) = Fx \oplus Fy \oplus Fz$ is the Lie algebra having the relations: $[x, y] = 0, [x, z] = ax, [y, z] = by$ with $a \neq b$.
- (2) If L has (L_1, L_{-1}) for a plus-minus pair, then L is isomorphic to either sl_2 or $\mathfrak{H}$.

Remark If $a = b$ is non-zero then we have $K(a, b) = K(a, a) \simeq K(1, 1)$ and this is nothing but our algebra $\mathfrak{g}$ of Theorem 2.3 which does not have a plus-minus pair.

Next we will briefly discuss isomorphism classes among the Lie algebras $K(a, b)$. Suppose that $K(a, b)$ is isomorphic to $K(a', b')$. If a or b is 0, then, by considering the derived subalgebra, we see that one of a' and b' is 0. Furthermore we see $K(0, c) \simeq K(c, 0) \simeq K(0, 1)$ for nonzero $c \in F$. Next we suppose that both a, b are nonzero. Then, we also see $K(a, b) \simeq K(a/b, 1)$. Suppose $\phi : K(a, 1) \rightarrow K(a', 1)$ is an isomorphism for nonzero $a, a' \in F$, where $K(a, 1)$ is generated by x, y, z with $[x, y] = 0, [x, z] = ax, [y, z] = y$, and $K(a', 1)$ is generated by x', y', z' with $[x', y'] = 0, [x', z'] = a'x', [y', z'] = y'$. Put $z'' = \phi(z) = a''x' + b''y' + c''z' \in K(a', 1)$. Then considering the derived algebras we find $\phi(Fx \oplus Fy) = Fx' \oplus Fy'$, and so we see that c'' is nonzero and that

$$\text{ad } z \mid_{Fx \oplus Fy} = \begin{pmatrix} -a & 0 \\ 0 & -1 \end{pmatrix}$$

must be similar to

$$\text{ad } z''|_{Fx' \oplus Fy'} = \begin{pmatrix} -c''a' & 0 \\ 0 & -c'' \end{pmatrix}.$$

Therefore, $a = c''a'$, $1 = c''$ or $a = c''$, $1 = c''a'$. This means $a' = a^{\pm 1}$. Conversely, we can easily confirm $K(a, 1) \simeq K(1/a, 1)$. We set $\mathfrak{P}(F) = \{ \{u, u^{-1}\} \mid u \in F, u \neq 0, 1 \} \cup \{ \{0\} \}$. Then, the isomorphism classes of the Lie algebras $K(a, b)$, having plus-minus pairs, are parametrized by the set $\mathfrak{P}(F)$. Here the isomorphism class of $K(0, a)$ corresponds to $\{0\}$ while that of $K(a, 1)$ to $\{a, a^{-1}\} = \{a^{-1}, a\}$ for $a \in F, a \neq 0$.

We next assume that F is an algebraically closed field of characteristic 0, and will study the three dimensional Lie algebras over F having a regular plus-minus pair. Let L be such an algebra and let (P, M) be a regular plus-minus pair of L . Then we can choose nonzero elements $x \in P$ and $y \in M$ as well as an involutive automorphism σ of L such that $[x, y] \neq 0$ and $\sigma(x) = y$. Put $z = [x, y]$, and set $u = x + y$ and $v = x - y$. Let $L_1 = Fu$ (the 1-eigenspace of σ) and $L_{-1} = Fv \oplus Fz$ (the -1 -eigenspace of σ). Then, $[u, v] = [x + y, x - y] = -2[x, y] = -2z$. We write $[z, x] = ax + by + cz$. Then we obtain $[z, y] = -\sigma([z, x]) = -\sigma(ax + by + cz) = -bx - ay + cz$. The Jacobi identity implies

$$[[x, y], z] + [[y, z], x] + [[z, x], y] = 0$$

and this gives $[bx + ay - cz, x] + [ax + by + cz, y] = 0$. Therefore,

$$\{-az - c(ax + by + cz)\} + \{az + c(-bx - ay + cz)\} = 0$$

and $c(a + b)x + c(a + b)y = 0$, which implies $c = 0$ or $a + b = 0$.

Case 1: $c = 0$.

In this case, we have $[x, y] = z$, $[z, x] = ax + by$, $[z, y] = -bx - ay$. We write the matrix of $\text{ad}(z)$ restricted to the space $Fx \oplus Fy$ as

$$\text{ad } z = \begin{pmatrix} a & -b \\ b & -a \end{pmatrix}.$$

Then, its characteristic polynomial is $t^2 - a^2 + b^2$. Hence, $\text{ad } z|_{Fx \oplus Fy}$ is similar to one of

$$\begin{pmatrix} \lambda & 0 \\ 0 & -\lambda \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},$$

where $\lambda = (a^2 - b^2)^{1/2}$. If $\lambda \neq 0$, that is $a^2 - b^2 \neq 0$, then we have certain elements $x', y' \in Fx \oplus Fy$ such that $[z', x'] = \lambda'x'$, $[z', y'] = -\lambda'y'$ with $z' = [x', y'] \neq 0$ and $\lambda' \neq 0$. This means that $L \simeq sl_2$. If $a = b = 0$, then we see $L \simeq \mathfrak{h}$. Now we suppose $a = b \neq 0$. Thus, we have $[z, u] = 0$, $[z, v] = 2au$, $[u, v] = -2z$. Put $\mu = (-a)^{1/4}$, and set $z' = z/\mu$, $u' = \mu u$, $v' = v/(2\mu^2)$. Then,

$$\begin{aligned} [v', u'] &= [v, u]/(2\mu) = z/\mu = z', \\ [v', z'] &= (-a)u/(\mu^3) = u', \\ [u', z'] &= 0. \end{aligned}$$

As is easy to check, one possible realization for this Lie algebra is obtained by taking

$$v' = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad u' = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad z' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

and

$$\sigma(v') = -v', \sigma(u') = u', \sigma(z') = -z'.$$

Next we suppose $a = -b \neq 0$. Thus, we have $[z, v] = 0, [z, u] = 2av, [u, v] = -2z$. Put $\nu = a^{1/4}$, and set $z' = z/\nu, u' = -u/(2\nu^2), v' = \nu v$. Then,

$$\begin{aligned} [u', v'] &= -[u, v]/(2\nu) = z/\nu = z', \\ [u', z'] &= -[u, z]/(2\nu^3) = \nu v = v', \\ [v', z'] &= 0. \end{aligned}$$

Here we can have the following realization:

$$u' = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad v' = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad z' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

and

$$\sigma(u') = u', \sigma(v') = -v', \sigma(z') = -z'.$$

Case 2: $c \neq 0, a + b = 0$.

In this case, we have $[x, y] = z, [z, x] = [z, y] = ax - ay + cz$, and hence $[u, v] = -2z, [z, u] = 2av + 2cz, [z, v] = 0$. Set $u' = -u/(2c), v' = cv, z' = z$. Then,

$$\begin{aligned} [u', v'] &= -[u, v]/2 = z = z', \\ [u', z'] &= -[u, z]/(2c) = av/c + z = av'/(c^2) + z, \\ [v', z'] &= 0. \end{aligned}$$

In this case we see the following gives a realization for our Lie algebra.

$$u' = \begin{pmatrix} 0 & r & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad v' = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad z' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

and

$$\sigma(u') = u', \sigma(v') = -v', \sigma(z') = -z'$$

with $r = a/(c^2)$.

In order to state a result we put $\mathbf{P}^1(F) = F \cup \{\infty\}$. For each $r \in \mathbf{P}^1(F)$, we define $\mathfrak{A}(r)$ to be the three dimensional Lie algebra spanned by $A(r), X, Y$, where

$$\begin{aligned} A(r) &= \begin{pmatrix} 0 & r & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (r \in F), \quad A(\infty) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ X &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

With this notation we see that the preceding arguments have proven the following result.

Theorem 2.5. Let F be an algebraically closed field of characteristic 0. Let L be a three dimensional Lie algebra with a regular plus-minus pair. Then, L is isomorphic to one of sl_2 , $\mathfrak{H}$ or $\mathfrak{A}(r)$ for $r \in \mathbf{P}^1(F)$.

We now discuss isomorphisms between these algebras. The dimension of the derived algebra of sl_2 is three, that of $\mathfrak{H}$ is one while that of $\mathfrak{A}(r)$, for any $r \in \mathbf{P}^1(F)$, is two so these algebras can never be isomorphic. Next we suppose $\mathfrak{A}(r) \simeq \mathfrak{A}(\infty)$ with $r \in F$. Then, as is seen by considering the action on the derived algebra which is spanned by X, Y we see both matrices

$$\begin{pmatrix} 0 & r \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad a \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

are similar for some nonzero $a \in F$. Hence, their characteristic polynomials $t(t-1)-r$ and t^2-a must coincide. This however is impossible so therefore, $\mathfrak{A}(r) \not\simeq \mathfrak{A}(\infty)$. Next we suppose $\mathfrak{A}(r) \simeq \mathfrak{A}(s)$ with $r, s \in F$. Then, both matrices

$$\begin{pmatrix} 0 & r \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad a \begin{pmatrix} 0 & s \\ 1 & 1 \end{pmatrix}$$

are similar for some nonzero $a \in F$. Hence, their characteristic polynomials $t(t-1)-r$ and $t(t-a)-as$ must coincide, which implies $a = 1$ and $r = s$. Therefore, $\mathfrak{A}(r) \not\simeq \mathfrak{A}(s)$ if $r \neq s$.

Finally we want to remark that the Lie algebra $K(u, 1)$ with $u \in F$ is isomorphic to $\mathfrak{A}(\infty)$ if $u = -1$, and is isomorphic to $\mathfrak{A}(-\frac{u}{(u+1)^2})$ if $u \neq -1$. This is because

$$\begin{pmatrix} -u & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

are similar if $u = -1$, and

$$\begin{pmatrix} -u & 0 \\ 0 & -1 \end{pmatrix} \quad \text{and} \quad a \begin{pmatrix} 0 & r \\ 1 & 1 \end{pmatrix}$$

with $a = -(u+1)$ and $r = -\frac{u}{(u+1)^2}$ are similar if $u \neq -1$. As a consequence of these computations, we see that a three dimensional three graded Lie algebra has a regular plus-minus pair or no plus-minus pair.

3 Plus-Minus pairs in some general classes of Lie algebras.

In this section we will show that Borchers Lie algebras have plus-minus pairs. Since these generalize the well-known Kac-Moody Lie algebras our results apply to these as well. We next go on to investigate $\mathbf{Z}^n$ -graded Lie algebras and see that with certain other assumptions these also have plus-minus pairs. Our method is to establish slightly more general factorization results in the universal enveloping algebras of these Lie algebras and then show how this gives rise to plus-minus pairs. The techniques are quite general and no doubt apply to other situations as well.

Let $\mathfrak{g}$ be a rank l Borcherds Lie algebra over F with the standard Cartan subalgebra $\mathfrak{h}$ and Chevalley generators $\{ e_1, \dots, e_l, f_1, \dots, f_l \}$, and let $\mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}]$ the derived subalgebra of $\mathfrak{g}$. Put $\mathfrak{h}' = \mathfrak{h} \cap \mathfrak{g}'$, and take a complement $\mathfrak{h}''$ of $\mathfrak{h}'$ in $\mathfrak{h}$ with $\mathfrak{h} = \mathfrak{h}'' \oplus \mathfrak{h}'$. Then, $\mathfrak{g} = \mathfrak{h}'' \oplus \mathfrak{g}'$. Let $\mathfrak{g}_+$ be the subalgebra of $\mathfrak{g}$ generated by $e_1, \dots, e_l$, and $\mathfrak{g}_-$ the subalgebra of $\mathfrak{g}$ generated by $f_1, \dots, f_l$ (cf. [4], [7], [8], [9], [10], [12]).

Proposition 3.1. Let $\mathfrak{g}$ be a rank l Borcherds Lie algebra, and let $I \cup J = \{ 1, 2, \dots, l \}$ be a partition of $\{ 1, 2, \dots, l \}$ into disjoint subsets. Then,

$$U(\mathfrak{g}) = \left(\prod_{i \in I} U(Fe_i) \right) U(\mathfrak{g}_-) U(\mathfrak{h}'') U(\mathfrak{g}_+) \left(\prod_{j \in J} U(Ff_j) \right).$$

Proof. Let $\mathfrak{g}_+^i$ be the standard homogeneous complementary subalgebra of Fe_i in $\mathfrak{g}_+$, and $\mathfrak{g}_-^i$ the standard homogeneous complementary subalgebra of Ff_i in $\mathfrak{g}_-$. For each $k = 1, \dots, l$ we put $h_k = [e_k, f_k]$ and $\mathfrak{h}_k = Fh_{k+1} \oplus \dots \oplus Fh_l$, and we set $I_k = I \cap \{ 1, \dots, k \}$ and $J_k = J \cap \{ 1, \dots, k \}$. We make free use of the PBW Theorem as well as the fact that $Fe_i \oplus Ff_i \oplus Fh_i$ is either sl_2 or $\mathfrak{h}$ so has a regular plus-minus pair.

If $1 \in I$, then

$$\begin{aligned} U(\mathfrak{g}) &= U(\mathfrak{g}_-) U(\mathfrak{h}) U(\mathfrak{g}_+) \\ &= U(\mathfrak{g}_-^1) U(Ff_1) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(Fh_1) U(Fe_1) U(\mathfrak{g}_+^1) \\ &= U(\mathfrak{g}_-^1) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(Ff_1) U(Fh_1) U(Fe_1) U(\mathfrak{g}_+^1) \\ &= U(\mathfrak{g}_-^1) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(Fe_1) U(Ff_1) U(Fe_1) U(\mathfrak{g}_+^1) \\ &= U(Fe_1) U(\mathfrak{g}_-^1) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(Ff_1) U(Fe_1) U(\mathfrak{g}_+^1) \\ &= U(Fe_1) U(\mathfrak{g}_-) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(\mathfrak{g}_+). \end{aligned}$$

In the other case when $1 \in J$ by using the same type of argument we have

$$U(\mathfrak{g}) = U(\mathfrak{g}_-) U(\mathfrak{h}'') U(\mathfrak{h}_1) U(\mathfrak{g}_+) U(Ff_1).$$

If we began with

$$U(\mathfrak{g}) = \left(\prod_{i \in I_k} U(Fe_i) \right) U(\mathfrak{g}_-) U(\mathfrak{h}'') U(\mathfrak{h}_k) U(\mathfrak{g}_+) \left(\prod_{j \in J_k} U(Ff_j) \right),$$

then, again using the same method, we can obtain

$$U(\mathfrak{g}) = \left(\prod_{i \in I_{k+1}} U(Fe_i) \right) U(\mathfrak{g}_-) U(\mathfrak{h}'') U(\mathfrak{h}_{k+1}) U(\mathfrak{g}_+) \left(\prod_{j \in J_{k+1}} U(Ff_j) \right).$$

Thus after several applications of this process we reach the stated result. Q.E.D.

We next see that this gives the desired plus-minus pair.

Corollary 3.2. Let $\mathfrak{g}$ be a Borcherds Lie algebra of finite rank. Then,

$$U(\mathfrak{g}) = U(\mathfrak{g}_{\pm}) U(\mathfrak{g}_{\mp}) U(\mathfrak{h}'') U(\mathfrak{g}_{\pm}).$$

Hence, perfect Borcherds Lie algebras have plus-minus pairs. In particular, perfect Kac-Moody Lie algebras or, more generally, perfect Brocherds Lie have regular plus-minus pairs.

Proof. We just take one of I and J to be empty. This leads to the result. Then, for example, let $P = \mathfrak{h}'' \oplus \mathfrak{g}_+$ and $M = \mathfrak{g}_-$. This gives a plus-minus pair. Q.E.D.

We next generalize the previous discussion by considering $\mathbf{Z}^n$ -graded Lie algebras. Thus, let $Q = \oplus_{i=1}^n \mathbf{Z}\alpha_i$ be a free abelian group of rank n generated by $\alpha_1, \dots, \alpha_n$, and let $\mathfrak{g} = \oplus_{\alpha \in Q} \mathfrak{g}_{\alpha}$ be a Lie algebra graded by Q . Put $\Delta = \{ \alpha \in Q \mid \mathfrak{g}_{\alpha} \neq 0 \}$. We also assume that $\mathbf{Z}\alpha_1 \cap \Delta = \{ 0, \pm\alpha_1 \}$, and that $L = \mathfrak{g}_{\alpha_1} \oplus [\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1}] \oplus \mathfrak{g}_{-\alpha_1}$ is a subalgebra with a plus-minus pair $(\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1})$ in L . (Thus, if L is three dimensional Proposition 2.4 implies L is isomorphic to either sl_2 or $\mathfrak{h}$.) We also suppose that there exists a complementary subalgebra $\mathfrak{g}'_0$ of $[\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1}]$ in $\mathfrak{g}_0$ with $\mathfrak{g}_0 = [\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1}] \oplus \mathfrak{g}'_0$. An element $\alpha = \sum_{i=1}^n c_i \alpha_i \in Q$ is called positive (resp. negative), that is $\alpha > 0$, if there is an index i satisfying $c_i > 0$ (resp. $c_i < 0$) and $c_{i+1} = c_{i+2} = \dots = c_n = 0$. Put $\Delta_+ = \{ \alpha \in \Delta \mid \alpha > 0 \}$ and $\Delta_- = \{ \alpha \in \Delta \mid \alpha < 0 \}$. Let $\mathfrak{g}_{\pm} = \oplus_{\alpha \in \Delta_{\pm}} \mathfrak{g}_{\alpha}$, and $\mathfrak{g}'_{\pm} = \oplus_{\alpha \in \Delta_{\pm} \setminus \{\alpha_1\}} \mathfrak{g}_{\alpha}$. Then, $\mathfrak{g}_{\pm} = \mathfrak{g}_{\pm\alpha_1} \oplus \mathfrak{g}'_{\pm}$, and we see that $\mathfrak{g}_{\pm\alpha_1} \oplus \mathfrak{g}'_{\mp}$ are subalgebras. In this situation we have the following result.

Proposition 3.3. Let $\mathfrak{g} = \oplus_{\alpha \in Q} \mathfrak{g}_{\alpha}$ be a graded Lie algebra with the extra conditions as above. Then,

$$U(\mathfrak{g}) = U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}_{-}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{+})$$

Moreover, letting $P = \mathfrak{g}_{+} \oplus \mathfrak{g}'_0$ and $M = \mathfrak{g}_{-}$ gives a plus-minus pair for $\mathfrak{g}$.

Proof. Using our assumptions we see that

$$\begin{aligned} U(\mathfrak{g}) &= U(\mathfrak{g}_{-}) U(\mathfrak{g}_0) U(\mathfrak{g}_{+}) \\ &= U(\mathfrak{g}'_{-}) U(\mathfrak{g}_{-\alpha_1}) U(\mathfrak{g}'_0) U([\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1}]) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{+}) \\ &= U(\mathfrak{g}'_{-}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{-\alpha_1}) U([\mathfrak{g}_{\alpha_1}, \mathfrak{g}_{-\alpha_1}]) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{+}) \\ &= U(\mathfrak{g}'_{-}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}_{-\alpha_1}) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{+}) \\ &= U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{-}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{-\alpha_1}) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{+}) \\ &= U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{-}) U(\mathfrak{g}_{-\alpha_1}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}'_{+}) \\ &= U(\mathfrak{g}_{\alpha_1}) U(\mathfrak{g}_{-}) U(\mathfrak{g}'_0) U(\mathfrak{g}_{+}). \end{aligned}$$

Q.E.D.

Remark It can be seen that many EALA's and some of the root-graded Lie algebras (cf. [1], [2], [3]) satisfy the hypothesis of Proposition 3.3.

References

- [1] B. N. Allison, S. Azam, S. Berman, Y. Gao and A. Pianzola, Extended affine Lie algebras and their root systems, *Memoirs AMS* 126 (No. 603), Providence, 1997.

- [2] B. N. Allison, G. Benkart and Y. Gao , Central extensions of Lie algebras graded by finite root systems, *Math. Ann.* 316 (2000), 499 – 527.
- [3] S. Berman and R. V. Moody , Lie algebras graded by finite root systems and the intersection matrix algebras of Slodowy, *Invent. Math.* 108 (1992), 323 – 347.
- [4] R. E. Borcherds , Generalized Kac-Moody algebras, *J. Algebra* 115 (1988), 501 – 512.
- [5] J. E. Humphreys , “Introduction to Lie algebras and representation theory”, GTM 9, Springer-Verlag, New York, 1972.
- [6] N. Jacobson , “Lie algebras”, Interscience, New York, 1962.
- [7] V. G. Kac , Simple irreducible graded Lie algebras of finite growth, *Math. USSR-Izv.* 2 (1968), 211 – 230.
- [8] V. G. Kac , “Infinite dimensional Lie algebras”, 3rd edition, Cambridge University Press, Cambridge, 1990.
- [9] R. V. Moody , A new class of Lie algebras, *J. Algebra* 10 (1968), 211 – 230.
- [10] R. V. Moody and A. Pianzola , “Lie algebras with triangular decompositions”, J. Wiley & Sons, New York, 1995.
- [11] J. Tits , Théorie des groupes, “Résumé des cours et travaux (1980 – 1981)”, pp. 75 – 87, Collège de France, Paris, 1981.
- [12] M. Wakimoto , “Infinite-dimensional Lie algebras”, Translations of Mathematical Monographs, Iwanami Series of Modern Mathematics, 2001.